


RESEARCH

Open Access



Sodium butyrate alleviates deoxynivalenol-induced hepatic cholesterol metabolic dysfunction via ROR γ -mediated histone acetylation modification in weaning piglets

Qiufang Zong^{1†}, Huan Qu^{1†}, Yahui Zhao¹, Haoyu Liu¹, Shenglong Wu^{1,2}, Shuai Wang³, Wenbin Bao^{1,2*} and Demin Cai^{1,2*} 

Abstract

Background: Cholesterol is an essential component of lipid rafts in cell plasma membrane, which exerts a hepatoprotective role against mycotoxin exposure in pigs, and cholesterol metabolism is vulnerable to epigenetic histone acetylation. Therefore, our present study aimed to investigate whether a histone deacetylase inhibitor (sodium butyrate [NaBu]) could protect the porcine liver from deoxynivalenol (DON) exposure by modulating cholesterol metabolism. Herein, we randomly divided 28 pigs into four groups, which were fed an uncontaminated basal diet, contaminated diet (4 mg DON/kg), uncontaminated diet supplemented with 0.2% NaBu or 4 mg/kg DON contaminated diet (4 mg DON/kg) supplemented with 0.2% NaBu for 28 d.

Results: We found that the serum alanine transaminase (ALT), aspartate transaminase (AST), and alkaline phosphatase (ALP) were all increased in pigs exposed to DON, indicative of significant liver injury. Furthermore, the cholesterol content in the serum of DON-exposed pigs was significantly reduced, compared to the healthy Vehicle group. Transcriptome analysis of porcine liver tissues revealed that the cholesterol homeostasis pathway was highly enriched due to DON exposure. In which we validated by qRT-PCR and western blotting that the cholesterol program was markedly activated. Importantly, NaBu effectively restored parameters associated with liver injury, along with the cholesterol content and the expression of key genes involved in the cholesterol biosynthesis pathway. Mechanistically, we performed a ChIP-seq analysis of H3K27ac and showed that NaBu strongly diminished DON-increased H3K27ac genome-wide enrichment. We further validated that the elevated H3K27ac and H3K9ac occupancies on cholesterol biosynthesis genes were both decreased by NaBu, as determined by ChIP-qPCR analysis. Notably, nuclear receptor ROR γ , a novel regulator of cholesterol biosynthesis, was found in the hyperacetylated regions. Again, a remarkable increase of ROR γ at both mRNA and protein levels in DON-exposed porcine livers was drastically reduced by NaBu. Consistent with ROR γ expression, NaBu also hindered ROR γ transcriptional binding enrichments on these activated cholesterol biosynthesis genes like *HMGCR*, *SQLE*, and *DHCR24*. Furthermore, we conducted an in vitro luciferase reporter assay to verify that porcine ROR γ directly bonds to the promoters of the above target genes.

[†]Qiufang Zong and Huan Qu contributed equally to this work.

*Correspondence: wbbao@yzu.edu.cn; demincai@yzu.edu.cn

¹ College of Animal Science and Technology, Yangzhou University, Yangzhou 225009, PR China

Full list of author information is available at the end of the article



Conclusions: Collectively, our results demonstrate the utility of the natural product NaBu as a potential anti-mycotoxin nutritional strategy for regulating cholesterol metabolism via ROR γ -mediated histone acetylation modification.

Keywords: Cholesterol biosynthesis, DON, Histone acetylation, ROR γ , Sodium butyrate

Background

Mycotoxins are secondary metabolites produced by filamentous fungi and can be ingested by animals and humans by accident with a high risk of acute or chronic toxicity [1, 2]. Worldwide, approximately 60%–80% of cereals are contaminated by various molds every year, thus directly threatening livestock and poultry production [3]. Deoxynivalenol (DON) is one of the most widely distributed mycotoxins. Its effect is often dose- and/or exposure time dependent, and can cause severe damage to the liver, gastrointestinal tract, and other metabolic and immunity-related organs [4]. As the central metabolic site, liver is responsible for detoxification following mycotoxin exposure. Given the fact that DON can hardly be removed 100% by the approaches of physical elimination, chemical degradation, and biodegradation, a small amount of accumulation can still cause liver damage in pigs [5]. Therefore, it is a burning question to explore the novel strategy against DON-induced toxicity in the porcine liver via endogenous approaches.

Notably, as the liquid-ordered microdomains in the plasma membrane, lipid rafts play a hepato-protective role by maintaining cellular membrane integrity to diminish mycotoxin-induced hepatocyte damage [6]. Owing to that lipid rafts are composed of cholesterol and sphingolipids, maintenance of cholesterol homeostasis is of great importance to ameliorate mycotoxin-induced liver injury as documented [7]. In contrast, mycotoxin exposure can elicit altered cholesterol production and dynamic balance, likely via the modulation of genes involved in cholesterol metabolic processes [8]. This suggests these genes are critical players enrolled in the prevention of mycotoxin damage. Notably, studies have revealed that cholesterol metabolic genes are susceptible to epigenetic regulation including DNA/RNA methylation and histone modifications [9]. Especially in the liver of pigs, histone acetylation exerts pivotal actions in the cholesterol metabolic gene programming [10]. Despite an increasing number of histone deacetylase (HDAC) modulators being widely studied to control acetylated events, sodium butyrate (NaBu), a natural and endogenous HDAC inhibitor, has attracted more attention in the last decade [11]. As expected, NaBu has been proven to mediate multiple biological reactions, including cholesterol metabolism, possibly via the regulation of histone acetylation [12]. Importantly, NaBu is thought to ameliorate the liver lesions triggered by DON in piglets

[13]. However, the underlying mechanisms have not been elucidated.

Tremendous research progress has indicated that various nuclear receptors (NRs), including ROR α/γ , REV-ERB α/β , LXRs, and PPARs, work as transcription factors (TFs) and play a cardinal role in controlling cholesterol metabolism by recruiting histone marks and co-factors [14, 15]. Notably, among multiple NRs, ROR γ has been found to specifically regulate cholesterol biosynthesis over the typical TF SREBP2 in porcine liver organoids. ROR γ activation by agonists or ectopic expression upregulates cholesterol biosynthetic gene expression via the hyper-enriched histone active mark H3K27ac [10]. It is elegantly proven that ROR γ directly binds to the genes involved in cholesterol biosynthesis using a genome-wide ChIP-seq analysis, and facilitates the histones' acetylation to enhance the cholesterol biosynthesis rate [16]. It is worth mentioning that ROR γ is one of the crucial drivers of this process in newborn piglets, in close association with mycotoxin-induced hepatic cholesterol re-programming and histone acetylation modifications containing H3K27ac [8]. In spite of that the expression and function of part of the NR members are found to be modified by NaBu, the direct evidence of the crosstalk between ROR γ and NaBu has not yet been reported. Thus, we herein hypothesized that ROR γ acts as a vital player enrolled in NaBu-alleviated liver injury in DON-exposed piglets via epigenetic regulation of cholesterol metabolism programming by histone acetylation.

Methods

Given that piglets are vulnerable to DON exposure, we generated a liver injury condition in piglets by dietary DON administration and then studied the hepatoprotective effects of NaBu. Cholesterol homeostasis was evaluated based on the total concentration in the liver, as well as using transcriptome analysis of hepatic cholesterol metabolism gene expression profiles. H3K27ac ChIP-seq analysis was used to examine genome-wide histone modification to verify ROR γ actions on cholesterol metabolism gene programming. ROR γ transcriptional/translational expression, and *in vitro* luciferase reporter assays were used to further validate the ROR γ targetable functions via directly binding fashions. Our present results probe the potential of NaBu as a *bona fide* agent for anti-mycotoxin through ROR γ -programmed epigenetic mechanisms of cholesterol metabolic genes.

Experimental animals and sampling

The animal study proposal was approved by the Institutional Animal Care and Use Committee (IACUC) of the Yangzhou University Animal Experiments Ethics Committee (permit number: SYXK (Su) IACUC 2012–0029). All experiments were conducted in accordance with the relevant guidelines and regulations. Twenty-eight barrows (28-day-old) were housed in an animal center, and randomly assigned to the following four groups: Vehicle (basal diet), DON (basal diet containing 4 mg DON/kg), NaBu (basal diet supplemented with 0.2% NaBu), and DON+NaBu (basal diet containing 4 mg DON/kg and 0.2% NaBu) (Fig. 1A). All of the pigs had ad libitum access to feed and water. The basal diet was formulated to satisfy the nutrient requirements of swine [17]. The DON feed was kindly provided by Huazhong Agricultural University [18], and the DON content was determined using an AgraQuant DON ELISA kit (Romer Labs, Singapore) according to the manufacturer's protocol. NaBu was purchased from Sigma-Aldrich (St. Louis, MO, USA) and added to the basal diet to form a NaBu content of 0.2%. After 28 d of feeding, blood samples were taken from all pigs via puncture of the jugular vein and centrifuged at 4000 r/min for 20 min to obtain serum. Thereafter pigs were euthanized with sodium pentobarbital, and liver samples were collected and immediately frozen in liquid nitrogen and preserved at -80°C .

Analysis of biochemical parameters

The concentrations of serum AST, ALT, ALP, CHO (cholesterol), TG (triglyceride), and TBA (total bile acids) were determined using commercial assay kits (Alovet,

Suzhou, China) according to the manufacturer's protocols using ALOVISION LIC200 (Alovet, Suzhou, China). Moreover, the protein of the liver was extracted, and the protein supernatant was collected for the determination of the biochemical index.

Quantitative real-time PCR (qRT-PCR)

Total RNA was isolated using TRIzol reagent (Takara, Dalian, China). After synthesis from the RNA using HiScript[®] II Q Select RT SuperMix (Vazyme, Nanjing, China), cDNA was subjected to qRT-PCR amplification using a StepOne Plus Real-Time Quantitative PCR System (Applied Biosystems, CA, USA). The primers were synthesized by Bioengineering (Shanghai, China). Primer information is shown in Table 1. The relative level of each transcript was normalized to that of *GAPDH* and analysed according to the $2^{-\Delta\Delta\text{Ct}}$ method [19].

RNA-seq and GSEA

Total RNA was extracted from the livers of pigs in the Vehicle, DON, NaBu, and DON+NaBu groups and prepared for RNA-seq library construction. PCR products were purified using an AMPure XP system (Beckman Coulter, CA, USA), and libraries were validated using an Agilent Bioanalyzer 2100 system (Agilent Technologies, CA, USA). Sequencing was performed on an Illumina HiSeq 2000 sequencer at BGI Tech (Wuhan, China). Reads were aligned to the genome assembly Sscrofa11.1 using TopHat2. Normalized gene expression was calculated using the HTSeq program. Differential expression analyses of DON vs. Vehicle and DON+NaBu vs. DON were performed




Table 1 Real-time PCR primer sequences

Genes	Primer sequences (5'→3')
ACAT2	F: TAATGATGGTCTGCTGCTGTGG R: GCTTGCTTTATTGCCGGGATTGG
HMGCS1	F: AAGCACAGCCACCGAGCATATTC R: ACCATCCCACCCACACTGAAG
HMGCR	F: TGTGATTGGAGTTGGCACCATGTC R: ACACGCAAGCTGGGAAGAAAGTC
MVK	F: GTTGCTCAAGTCCTGCTGGTGTC R: AGGCTCACTTCCCACTGTTGTG
PMVK	F: GGTGGATGATGCTGAGTCAGAGTG R: GTGCTGCTCATCTCCGTGGTTC
MVD	F: GCCACCTGCTGGACACCTTC R: GGCGAAGATCACGGCGTTGG
FDFT1	F: GCGTCCACCCTCCTCACTCC R: CCCACACAGCCAGAGCCAAAG
SQLE	F: TGTGGACCTTTCTCGGCATTGC R: TAGCGACAGCGGTAGGACAGC
LSS	F: GAGGACCCGCTGGTCCA R: CCACACTGTTCTGTGCGC
NSDHL	F: TTTGTGATCGGGAACGGGAAGAAC R: TTCGTATTGGTGATGTGGAAGGC
DHCR24	F: CCTTCTCTCTGCCGCTCTC R: TGCCCTGCTCCTCCATTCCC
RORC	F: CAATGGAAGTGGTGTGGTCAGG R: GGGAGCGGGAGAAAGTCAAAGATG
GAPDH	F: ACATCATCCCTGCTTCTACTGG R: CTCGGACGCTGCTTCC

using DESeq of the R package. Genes with an adjusted P value < 0.05 and $|\log_2(\text{fold change})| > 1$ were defined as differentially expressed. Gene set enrichment analysis (GSEA 4.1.0) was applied to rank differentially expressed genes and enrich the biological processes and pathways.

Western blotting

Frozen liver tissue was lysed in RIPA buffer (Beyotime, Shanghai, China) containing protease inhibitors. After centrifugation at $12,000 \times g$ for 15 min at 4°C , the supernatant was obtained, and proteins were quantified using an Enhanced BCA Protein Assay Kit (Beyotime, Shanghai, China). Proteins were electrophoresed and electrotransferred onto PVDF membranes (Millipore, MA, USA). The membranes were blocked with 5% skimmed milk at room temperature (RT) for 2 h and incubated with primary antibodies at 4°C overnight. After washing 3 times with cold PBS and incubating with a secondary antibody at RT for 2 h, the membranes were visualised using an ECL detection reagent and a FluorChem FC3 system (ProteinSimple, CA, USA). The antibodies used are shown in Table 2.

Table 2 Western blotting antibodies

Antibody	Manufacturer	Catalogue number	Dilution
MVK	Santa-Cruz	sc-390669	1:1000
MVD	Santa-Cruz	sc-376975	1:1000
FDFT1	Santa-Cruz	sc-271602	1:1000
SQLE	Santa-Cruz	sc-271651	1:1000
HMGCS1	Santa-Cruz	sc-166763	1:1000
ROR γ	Invitrogen	14-6988-82	1:1000
GAPDH	Proteintech	10494-1-AP	1:5000
Anti-Mouse IgG	HuaBio	HA1006	1:10000
Anti-Rabbit IgG	HuaBio	HA1001	1:10000

ChIP-qPCR and ChIP-seq data analysis

Porcine livers from the Vehicle, DON, NaBu, and DON+NaBu groups were ground in liquid nitrogen and subjected to crosslinking in 1% formaldehyde for 6 min, followed by quenching with glycine for 6 min. After 3 washes with PBS, the pellets were collected by centrifugation and resuspended in lysis buffer (50 mmol/L HEPES-KOH, 140 mmol/L NaCl, 10% glycerol, 1 mmol/L EDTA, 0.25% Triton-X100 and 0.5% NP-40). The pellets were resuspended in washing buffer (10 mmol/L Tris, 1 mmol/L EDTA, 0.5 mmol/L EGTA and 200 mmol/L NaCl). The supernatants were discarded, and then the pellets were resuspended in a shearing buffer (0.1% SDS, 1 mmol/L EDTA and 10 mmol/L Tris-HCl) for further sonication using a Covaris M220 (MA, USA) according to the manufacturer's recommendations. Crude chromatin fragments were precipitated using antibodies and Protein G beads. After washing and adding proteinase K and RNase A, purified ChIP DNA was collected for further ChIP-seq and ChIP-qPCR assays.

The antibodies used in the ChIP assay were anti-ROR γ serum (generated by Covance), H3K9ac (Abcam, ab4441), and H3K27ac (Abcam, ab4729). Collected DNA was quantified using an Agilent Bioanalyzer 2100 for sequencing on an Illumina HiSeq 2000 Sequencer (BGI, Wuhan, China). Raw single-end sequencing data (ChIP-sequencing reads) were checked for quality using FastQC (v0.11.9). The data were then aligned to the Sscrofa11.1 reference genome using Bowtie 1.2.3, followed by peak calling using MACS2 (2.1.1). Uniquely mapped tag filtering and deduping were used for peak calling by model-based analysis for ChIP-seq (MACS; 2.1.0) to identify regions of ChIP-seq enrichment over the background. Normalized genome-wide signal-coverage tracks from raw read alignment files were built using MACS2 and bedTools (<https://github.com/arq5x/bedtools2>). ChIP-seq signals at enriched genomic

regions (avgprofile and heatmap) were visualised using deepTools (<https://deeptools.readthedocs.io/en/develop/index.html>). Peaks located in the promoter, 5'-UTR, 3'-UTR, exons, introns, and intergenic regions were annotated. Differential peaks in variant groups were picked using IGV software. The primers were designed based on the sequences of the peaks. ChIP-qPCR assays were performed according to the manufacturer's recommendations. The primers are shown in Table 3.

Vector constructs and dual-luciferase reporter assay

For cholesterol homeostasis gene vector construction, we synthesized fragments of the *HMGCR*, *SQLE*, and *DHCR24* promoters with or without mutation of the putative RORC binding sites. *HMGCR*-wt and *HMGCR*-mut vectors were constructed by inserting the synthesized fragments of the *HMGCR* promoters from Chr2: 84,370,135–84,370,376 (Sscrofa11.1) and the mutant form from AGGTCA to CCCAAC into the pGL3-basic luciferase reporter vector. Similarly, *SQLE*-wt and *SQLE*-mut vectors were constructed by inserting the synthesized fragments of the *SQLE* promoters from Chr4: 14,684,789–14,685,079 (Sscrofa11.1) and the mutant form from TGACCT to GTTGGG into the pGL3-basic luciferase reporter vector. *DHCR24*-wt and *DHCR24*-mut vectors were constructed by inserting the synthesized fragments of the *DHCR24* promoters from Chr6: 157,469,893–157,470,158 (Sscrofa11.1) and the mutant form from TGACCT to GTTGGG into the pGL3-basic luciferase reporter vector. After 293T cells reached 70% confluence in 12-well plates, the wild types or mutant types of *HMGCR*, *SQLE*, and *DHCR24* genes were co-transfected with the *RORC* overexpression (*RORC*) vector. The renilla plasmid was co-transfected for normalization. Fluorescence activity was detected 36 h later using a dual-luciferase reporter system (Promega Corporation, WI, USA) according to the manufacturer's instructions. Data were obtained from 3 independent experiments, each conducted in sextuplicate.

Table 3 ChIP-qPCR primer sequences

Gene	Primer sequences (5'→3')
<i>HMGCR</i>	F: TCTGAAGAAGTTTAAGGGAA R: GCTTGGGTGCTGTG
<i>SQLE</i>	F: CAGGACTGGCTTCTTC R: ATGGGTTCCGGCACTAG
<i>HSD17B7</i>	F: ATGCTTGATAGGGTTT R: CAGAGGGTGTGAGTTT
<i>DHCR24</i>	F: ATTGCTGTGGCGTAGA R: GGTGCTTCAGAGGAGG
<i>LDLR</i>	F: ACAGCCATAGCAATACA R: TCAATGAGGAGGTAGGT

Bioinformatic analysis using a clinical dataset

Data for mRNA expression-based liver cancer were downloaded from UCSC Xena datasets (<https://xenabrowser.net/hub/>). The data were then log2 transformed and quantile normalized before further analysis. Pearson's correlation metric was computed between each gene using the 'cor' function in R, and the 'ggplot2' R package was used for further visualisation.

Statistical analysis

All data were analysed using SPSS Statistics 21 software. Data are presented as the means ± SEM. Statistical analysis was performed using one-way ANOVA to compare the means. $P < 0.05$ was considered significant.


Results

Sodium butyrate alleviates DON-induced liver injury in piglets

To verify that DON can cause liver injury in piglets, the key parameters of liver injury including ALT, AST, and ALP levels in the serum and liver were analysed. As expected, elevated levels of ALT, AST, and ALP in the serum were observed in the DON-exposed piglets while these were recovered by NaBu treatment. Even though serum ALT and AST levels are increased in NaBu group, the effect is not significant and without statistical significance (Fig. 1B). Consistent with the results in serum, although the ALT and AST levels were not different among these four groups, NaBu markedly reduced the contents of ALP in the DON-exposed piglets compared to DON treatment alone (Fig. 1C).

Sodium butyrate prevents DON-caused cholesterol metabolic abnormality

To explore the core transcriptional program in the liver of piglets upon DON contamination with or without NaBu, we performed RNA-seq analysis using liver tissues from the Vehicle, DON, NaBu, and DON+NaBu groups. We performed PCA by comparing gene expression levels using count data from the four groups. The results showed that single DON or NaBu treatment caused the obvious difference of the genes compared with Vehicle, while the combined DON with NaBu approach promoted the recovery of part of DON-altered genes (Fig. 2A). One thousand three hundred thirty-nine differentially expressed genes were identified between Vehicle and DON groups, among which were 793 up-regulated and 546 down-regulated genes by DON compared to that of Vehicle. Moreover, 269 differentially expressed genes were identified between DON and DON+NaBu groups, in which NaBu supplementation induced the upregulation of 188 and downregulation of 81 genes compared




with DON treatment alone (Fig. 2B). Venn diagram manifested that the expression of 788 upregulated genes in the DON-treated piglets was inhibited by NaBu. In contrast, NaBu enhanced the expression of 235 genes in the porcine liver that was suppressed by DON exposure alone (Fig. 2C). Clustering of genes with expression significantly altered by DON (relative to Vehicle) showed a high degree of concordance in expression changes induced by NaBu (Fig. 2D). In line with these results, GSEA analysis revealed that the cholesterol homeostasis pathway was highly enriched and inhibited in the DON+NaBu groups versus DON treatment (Fig. 2E). Again, a pathway-focused analysis showed that the majority of cholesterol biosynthesis genes were significantly up-regulated by DON and then were decreased with the treatment of NaBu with DON (Fig. 2F).

To validate the RNA-seq results, the related mRNA expression was further measured by qRT-PCR. The expression of the key genes including *ACAT2*, *HMGCS1*, *HMGCR*, *MVK*, *PMVK*, *MVD*, *FDFT1*, *SQLE*, *LSS*, *NSDHL*, and *DHCR24* involved in cholesterol biosynthesis was significantly up-regulated by DON exposure, and was strongly inhibited in the group with the treatment of DON+NaBu compared to that of DON alone (Fig. 3A). Similarly, the immunoblotting results displayed that NaBu supplementation efficiently diminished the HMGCS1 and FDFT1 proteins expression to the comparable level of vehicle in the DON-exposed piglets, compared to that of single DON treatment (Fig. 3B and C).

This was in association with the contents of CHO, TG, and TBA in the porcine serum with indicated treatment, in which the contents of the aforementioned index were significantly reduced in the DON-exposed group while increased in combination with NaBu (Fig. 3D-F).

NaBu supplementation abolishes the enhanced genome-wide H3K27ac occupancy on cholesterol biosynthesis genes

To further explore the underlying mechanism by which NaBu regulates the cholesterol biosynthesis pathway, ChIP-seq analysis of histone acetylated mark H3K27ac was performed. We first demonstrated that the location of the binding sites was drastically shifted to promoter regions treated with DON, whereas NaBu markedly blunted the shift to display the similar pattern of the located binding regions to Vehicle (Fig. 4A). Moreover, DON exposure dramatically increased the genome-wide H3K27ac association with its targets, which was reduced by NaBu supplementation (Fig. 4B). Concomitant with the up-regulated transcripts involved in cholesterol metabolism, hyper-enriched H3K27ac was observed at the cholesterol metabolic gene loci in the DON-exposed piglets. Obviously, NaBu remarkably decreased these relative enrichments as shown in the plots of Fig. 4C. Notably in the cholesterol biosynthetic program, as shown in the heat-map of H3K27ac signal intensity (Fig. 4D), NaBu supplementation caused the alterations of H3K27ac occupancies were in agreement with the



mRNA levels of the aforementioned cholesterol biosynthesis pathway.

Histone acetylated marks H3K27ac and H3K9ac contribute to NaBu-modulated key cholesterol biosynthetic genes

Having revealed the crucial functions of histone acetylation in the regulation of the cholesterol metabolic pathway, we next examined which genes were susceptible to DON or NaBu exposure. Combined with the up-regulated transcriptional expression, H3K27ac enrichment showed a dramatic increment at the enhancers of *HMGCR*, *SQLE*, *HSD17B7*, *DHCR24* and *LDLR* in response to DON exposure, the significant loss caused by NaBu was also seen in the signal visualisation (Fig. 5A). Importantly, in addition to H3K27ac, we performed ChIP-qPCR analysis to validate and quantify the observations of histone acetylation. As shown in Fig. 5B and C, the cholesterol biosynthesis genes all displayed higher enrichments on the specific binding sites in the DON groups and these were diminished with NaBu treatment.

ROR γ is a key player involved in histone acetylation

Nuclear receptor ROR γ has been found to play a crucial role in the process of cholesterol biosynthesis [8]. Therefore, we aimed to further determine whether

NaBu-regulated cholesterol biosynthesis is also reprogrammed by ROR γ . Clinical dataset indicated a strongly positive correlation between the expression of *RORC* and *HMGCR* ($r = 0.34$, $P < 0.001$), *SQLE* ($r = 0.2$, $P < 0.001$), *DHCR24* ($r = 0.18$, $P < 0.001$), *LDLR* ($r = 0.33$, $P < 0.001$), *HMGCS1* ($r = 0.27$, $P < 0.001$), *FDFT1* ($r = 0.25$, $P < 0.001$) and *NSDHL* ($r = 0.22$, $P < 0.001$) (Fig. 6A). We then evaluated the FPKM value and mRNA expression of *RORC* gene, and the results revealed that the increased *RORC* expression in DON-exposed porcine livers was dramatically reduced by NaBu (Fig. 6B and C). In line with the mRNA expression results, NaBu supplementation alleviated DON-induced upregulation of ROR γ protein expression (Fig. 6D and E). To determine whether ROR γ is involved in histone modification in the present study, we performed ChIP-qPCR to detect the ROR γ occupancy in the region of H3K27ac enrichment. As shown in Fig. 7A, consistent with the ROR γ expression and histone acetylation modification results, NaBu hindered ROR γ transcriptional binding enrichment on activated cholesterol biosynthesis genes. Moreover, in a dual-luciferase reporter assay, we found that promoters or enhancers of *HMGCR*, *SQLE*, and *DHCR24* genes were highly responsive to ROR γ -dependent activation.





Fig. 4 NaBu supplementation abolishes the enhanced genome-wide H3K27ac occupancy on cholesterol biosynthesis genes. **A** H3K27ac enrichment regions were localized within a 3-kb region of the promoter, exon, intron, UTR, and intergenic regions in porcine liver. **B** ChIP-seq profiles of H3K27ac signal intensity within ± 3 -kb windows around the center of peak regions. **C** ChIP-seq profiles of H3K27ac binding on genes involved in the cholesterol biosynthesis pathway. **D** Heatmaps of ChIP-seq signal intensities of genes in the cholesterol biosynthesis pathway

However, the mutated type of the putative RORy binding domain diminished the RORy-mediated activation (Fig. 7B).

A graphical illustration of the mechanism by which NaBu alleviates DON-induced disturbances in cholesterol biosynthesis via RORy-mediated histone acetylation is shown in Fig. 8. Our findings indicate that DON exposure causes liver injury, resulting in a decrease in cholesterol content in the serum and activation of the RORy-mediated cholesterol biosynthesis pathway. When cholesterol levels in the body are elevated, RORy expression and the continuous synthesis of cholesterol are in turn inhibited. This process is regulated by H3K27ac and H3K9ac acetylation.

Discussion


Humans and animals are vulnerable to mycotoxin exposure because of the high toxicity and absorbability. In pigs, in particular, the absorption rate of DON is up to 66% as documented [20], which causes severe consequences in nearly all organs. Similar to previous findings of liver lesions induced by DON [21], here, we also found the obviously increased liver injury parameters in blood and liver, as ALT, AST and ALP were at pathological levels in the DON-treated piglets. It has been suggested that lipid dysfunction is closely associated with mycotoxin-induced liver lesions [22]. Notably, lipid rafts play cellular-protection roles in maintaining membrane properties and H^+ -ATPase activity against




mycotoxin-induced structural failure of cell membrane by activating ceramide synthesis at the endoplasmic reticulum [23]. Lipid rafts are sub-domains of the plasma membrane and consist of cholesterol and sphingolipids, thus cholesterol homeostasis was our primary concern in the present study. As speculated, we observed that the cholesterol content in the serum of piglets was significantly reduced by DON exposure, suggesting that the hepatocyte damage was attributed to cholesterol depletion at cell membranes. Although the potential anti-mycotoxin action of NaBu has been suggested previously [13], we were still excited that NaBu strongly restored the cholesterol content and key parameters involved in liver injury. Notably, in the transcriptome analysis, the whole cholesterol metabolism pathway in the DON group is dramatically driven to the comparable level of Vehicle when NaBu was supplemented.

Given the classic function of NaBu in histone acetylation, we reasonably evaluated histone modifications on the key genes involved in cholesterol metabolism using our present model. Indeed, we have previously demonstrated that mycotoxins from commercial farms, including DON, ochratoxin, zearalenone, and aflatoxin B1 could markedly decrease cholesterol levels in pigs via epigenetic histone modification [8]. In this study, we found that the cholesterol biosynthesis pathway was upregulated in porcine livers when piglets were supplemented with DON alone. Indeed, hepatic cholesterol biosynthesis is typically negatively regulated via a signal derived

from circular cholesterol loss. Because of the central site for cholesterol metabolism, this feedback loop facilitates the cholesterol balance among organs [24]. NaBu has been reported to modulate the expression of the genes involved in cholesterol biosynthesis and uptake [25]. Obviously, the increased expression of cholesterol biosynthesis genes in DON-treated piglets was down-regulated by NaBu supplementation. However, NaBu supplementation alone did not affect the cholesterol content and biosynthesis pathway, suggesting that a dose of 0.2% is most appropriate for anti-DON treatment in pigs, with no obvious side effects. Additional evidence of the safety of NaBu is the absence of an effect on ALT, AST, and ALP in the liver and serum compared with Vehicle group. Even though serum ALT and AST levels are increased in NaBu group. However, the effect is not significant and without statistical significance, and its levels are still within the normal physiological range [26–28]. Moreover, we also found that the genome-wide binding of histone acetylation mark H3K27ac was not changed by NaBu administration alone using ChIP-seq analysis. However, NaBu administration efficiently diminished DON-induced hyperacetylation both at the genome-wide level and of cholesterol synthesis genes. Indeed, NaBu, as a histone deacetylase inhibitor (HDACi), has been well documented to exert theoretical activity in epigenetically silenced genes by enhancing global histone acetylation [29]. Nevertheless, it is worth mentioning that HDACi has also been demonstrated to cause a similar amount






or even more genes which are decreased than elevated [30]. Importantly, the majority of these alterations in gene expression with an even larger fraction triggered by HDACi are shown to be down-regulated [31, 32]. Combining our present results with previous reports, NaBu exhibited a couple of genomic characteristics on the DNA-bound histone acetylation status. Firstly, NaBu is more likely to create targets for histone deacetylation around the TSS regions while these genes are frequently down-regulated. In contrast, NaBu-induced genome-wide histone hyperacetylation continually occurs along the nuclear periphery. Secondly, the hyperacetylation of specific genes caused by NaBu is transient, whereas the histone acetylation status dynamically changes in a time-dependent manner. Additionally, NaBu would reset the structures and functions to regulate gene expression when initiating histone deacetylation [33].

Another novel finding in our present study is the NR RORγ action that mediates the NaBu-reprogrammed cholesterol biosynthesis transcript. Recent studies have provided a number of clues to control cholesterol

metabolism in different scenarios by NRs including RORα/γ, REV-ERBα, and PPARα, directly or in combination with the classic cholesterol transcription factors SREBP2 and LXRs [14, 15, 34]. Importantly, RORγ has been demonstrated to drive cholesterol biosynthesis over SREBP2 by recruiting H3K27ac in a breast cancer cell line [16]. Notably, in a liver organoid model derived from piglets, RORγ was selected among a series of NRs as specifically modulating the cholesterol biosynthesis program [10]. In addition, we previously revealed that the hepatic protein content is positively correlated with the genome-wide binding enrichment of RORγ in mycotoxin-exposed piglets [8]. Herein, we further confirmed the potential regulatory effect of RORγ on the expression of cholesterol-related genes in pigs through a co-relationship analysis using clinical datasets and a luciferase reporter assay specifically targeting *HMGCGR*, *SQLE* and *DHCR24*. In spite of that NaBu is suggested to activate NRs such as PPARs and VDR as a metabolite [35–37], the reports of its direct effects on RORγ are lacking. Interestingly, RORγt, a thymus-specific



isoform of ROR γ , was shown to be inhibited by NaBu in Th17 cells [38]. In agreement, NaBu supplementation dramatically suppressed ROR γ expression at both the transcriptional and translational levels when exposed to DON. This inhibition was also shared at the chromatin binding occupancies that ROR γ enrichments were similar to those of Vehicle in the pigs supplemented with DON along with NaBu. However, it is still unclear whether NaBu inactivates ROR γ by directly binding to the ligand domain of ROR γ . This is a limitation of the present study and needs to be explored in further investigations.

Conclusions

Our study identified a novel mechanism in which ROR γ targets cholesterol biosynthesis by regulating histone acetylation. These results contribute to the understanding of the role of ROR γ in mycotoxin-induced liver injury in pigs. This study also illustrates a novel potential approach for mycotoxin poisoning prevention and intervention from the perspective of epigenetic modifications.

Abbreviations

NaBu: Sodium butyrate; DON: Deoxynivalenol; ALT: Alanine transaminase; AST: Aspartate transaminase; ALP: Alkaline phosphatase; HDAC: Histone deacetylase; NRs: Nuclear receptors; TFs: Transcription factors; CHO: Cholesterol; TG: Triglyceride; TBA: Total bile acids; qRT-PCR: Quantitative real-time PCR; RT: Room temperature; HDACi: Histone deacetylase inhibitor.

Acknowledgements

Not applicable.

Authors' contributions

QFZ and HQ performed most of the experiments. YHZ, HYL, SLW and SW participated in the experiments. DMC and WBB conceived the study. QFZ, HQ and DMC participated in study design and coordination and wrote the manuscript. All authors read and approved the final manuscript.

Funding

This study was supported by Key Research and Development Project (Modern Agriculture) of Jiangsu Province (BE2019341), Jiangsu Agricultural Science and Technology Innovation Fund (CX [20]2003), and the Priority Academic Program Development of Jiangsu Higher Education Institutions.

Availability of data and materials

The datasets analysed during the current study are available from the corresponding author upon request.

Declarations

Ethics approval and consent to participate

The animal study proposal was approved by the Institutional Animal Care and Use Committee (IACUC) of the Yangzhou University Animal Experiments Ethics Committee (permit number: SYXK (Su) IACUC 2012–0029). All experiments were conducted in accordance with the relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹College of Animal Science and Technology, Yangzhou University, Yangzhou 225009, PR China. ²Joint International Research Laboratory of Agriculture & Agri-Product Safety, Yangzhou University, Yangzhou 225009, PR China. ³Department of Animal

Nutrition and Feed Science, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan 430070, PR China.

Received: 17 May 2022 Accepted: 6 October 2022
Published online: 23 December 2022

References

- Tesfamariam K, De Boevre M, Kolsteren P, Belachew T, Mesfin A, De Saeger S, et al. Dietary mycotoxins exposure and child growth, immune system, morbidity, and mortality: a systematic literature review. *Crit Rev Food Sci Nutr*. 2020;60(19):3321–41. <https://doi.org/10.1080/10408398.2019.1685455>.
- Haque MA, Wang YH, Shen ZQ, Li XH, Saleemi MK, He C. Mycotoxin contamination and control strategy in human, domestic animal and poultry: a review. *Microb Pathog*. 2020;142:104095. <https://doi.org/10.1016/j.micpath.2020.104095>.
- Eskola M, Kos G, Elliott CT, Hajšlová J, Mayar S, Krška R. Worldwide contamination of food-crops with mycotoxins: validity of the widely cited 'FAO estimate' of 25. *Crit Rev Food Sci Nutr*. 2020;60(16):2773–89. <https://doi.org/10.1080/10408398.2019.1658570>.
- Pestka JJ. Deoxynivalenol: mechanisms of action, human exposure, and toxicological relevance. *Arch Toxicol*. 2010;84:663–79. <https://doi.org/10.1007/s00204-010-0579-8>.
- Yao Y, Long M. The biological detoxification of deoxynivalenol: a review. *Food Chem Toxicol*. 2020;145:111649. <https://doi.org/10.1016/j.fct.2020.111649>.
- Zhang Y, Qi XZ, Zheng JJ, Luo YB, Zhao CH, Hao JR, et al. Lipid rafts disruption increases ochratoxin A cytotoxicity to hepatocytes. *J Biochem Mol Toxicol*. 2016;30(2):71–9. <https://doi.org/10.1002/jbt.21738>.
- Adimulam T, Abdul NS, Chuturgoon AA. HepG2 liver cells treated with fumonisin B1 in galactose supplemented media have altered expression of genes and proteins known to regulate cholesterol flux. *World Mycotoxin J*. 2021;15(3):313–24. <https://doi.org/10.3920/WMJ2021.2723>.
- Li KQ, Li H, Zhang KX, Zhang JY, Hu P, Li YW, et al. Orphan nuclear receptor ROR γ modulates the genome-wide binding of the cholesterol metabolic genes during mycotoxin-induced liver injury. *Nutrients*. 2021;13(8):2539. <https://doi.org/10.3390/nu13082539>.
- Ferrari A, Fiorino E, Giudici M, Gilardi F, Galmozzi A, Mitro N, et al. Linking epigenetics to lipid metabolism: focus on histone deacetylases. *Mol Membr Biol*. 2012;29(7):257–66. <https://doi.org/10.3109/09687688.2012.729094>.
- Zhang KX, Li H, Xin ZM, Li YW, Wang XL, Hu Y, et al. Time-restricted feeding downregulates cholesterol biosynthesis program via ROR γ -mediated chromatin modification in porcine liver organoids. *J Anim Sci Biotechnol*. 2020;11:106. <https://doi.org/10.1186/s40104-020-00511-9>.
- Jiang LQ, Wang JJ, Liu ZY, Jiang AM, Li SQ, Wu D, et al. Sodium butyrate alleviates lipopolysaccharide-induced inflammatory responses by down-regulation of NF- κ B, NLRP3 signaling pathway, and activating histone acetylation in bovine macrophages. *Front Vet Sci*. 2020;7:579674. <https://doi.org/10.3389/fvets.2020.579674>.
- Nunes MJ, Milagre I, Schnekenburger M, Gama MJ, Diederich M, Rodrigues E. Sp proteins play a critical role in histone deacetylase inhibitor-mediated derepression of CYP46A1 gene transcription. *J Neurochem*. 2010;113(2):418–31. <https://doi.org/10.1111/j.1471-4159.2010.06612.x>.
- Wang S, Zhang C, Yang JC, Wang X, Wu KT, Zhang BY, et al. Sodium butyrate protects the intestinal barrier by modulating intestinal host defense peptide expression and gut microbiota after a challenge with deoxynivalenol in weaned piglets. *J Agric Food Chem*. 2020;68(15):4515–27. <https://doi.org/10.1021/acs.jafc.0c00791>.
- Wang B, Tontonoz P. Liver X receptors in lipid signalling and membrane homeostasis. *Nat Rev Endocrinol*. 2018;14(8):452–63. <https://doi.org/10.1038/s41574-018-0037-x>.
- Kojetin DJ, Burris TP. REV-ERB and ROR nuclear receptors as drug targets. *Nat Rev Drug Discov*. 2014;13(3):197–216. <https://doi.org/10.1038/nrd4100>.
- Cai DM, Wang JJ, Gao B, Li J, Wu F, Zou JX, et al. ROR γ is a targetable master regulator of cholesterol biosynthesis in a cancer subtype. *Nat Commun*. 2019;10(1):4621. <https://doi.org/10.1038/s41467-019-12529-3>.
- United States National Research Council. Nutrient requirements of swine. 11th ed. Washington: National Academy Press; 2012.
- Wang S, Yang JC, Zhang BY, Wu KT, Yang A, Li C, et al. Deoxynivalenol impairs porcine intestinal host defense peptide expression in weaned piglets and IPEC-J2 cells. *Toxins (Basel)*. 2018;10(12):541. <https://doi.org/10.3390/toxins10120541>.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{- $\Delta\Delta$ CT} method. *Methods*. 2001;25:402–8. <https://doi.org/10.1006/meth.2001.1262>.
- Goyarts T, Dänicke S. Bioavailability of the Fusarium toxin deoxynivalenol (DON) from naturally contaminated wheat for the pig. *Toxicol Lett*. 2006;163(3):171–82. <https://doi.org/10.1016/j.toxlet.2005.10.007>.
- Bai YS, Ma KD, Li JB, Li JP, Bi CP, Shan AS. Deoxynivalenol exposure induces liver damage in mice: inflammation and immune responses, oxidative stress, and protective effects of *Lactobacillus rhamnosus* GG. *Food Chem Toxicol*. 2021;156:112514. <https://doi.org/10.1016/j.fct.2021.112514>.
- Martins IJ. Overnutrition determines LPS regulation of mycotoxin induced neurotoxicity in neurodegenerative diseases. *Int J Mol Sci*. 2015;16(12):29554–73. <https://doi.org/10.3390/ijms161226190>.
- Gutiérrez-Nájera NA, Saucedo-García M, Noyola-Martínez L, Vázquez-Vázquez C, Palacios-Bahena S, Carmona-Salazar L, et al. Sphingolipid effects on the plasma membrane produced by addition of fumonisin B1 to maize embryos. *Plants (Basel)*. 2020;9(2):150. <https://doi.org/10.3390/plants9020150>.
- Ru P, Guo D. microRNA-29 mediates a novel negative feedback loop to regulate SCAP/SREBP-1 and lipid metabolism. *RNA Dis*. 2017;4(1):e1525. <https://doi.org/10.14800/rd.1525>.
- Nunes MJ, Moutinho M, Gama MJ, Rodrigues CM, Rodrigues E. Histone deacetylase inhibition decreases cholesterol levels in neuronal cells by modulating key genes in cholesterol synthesis, uptake and efflux. *PLoS One*. 2013;8(1):e53394. <https://doi.org/10.1371/journal.pone.0053394>.
- Wang PZ, Lu Z, He M, Shi BM, Lei XG, Shan AS. The effects of endoplasmic reticulum-resident selenoproteins in a nonalcoholic fatty liver disease pig model induced by a high-fat diet. *Nutrients*. 2020;12(3):692. <https://doi.org/10.3390/nu12030692>.
- Zhao DD, Wang JJ, Fan JJ, Fu Y, Ma YD, Li C, et al. Taurine prevents LPS-induced liver injury in weaned piglets. *Adv Exp Med Biol*. 2022;1370:497–505. https://doi.org/10.1007/978-3-030-93337-1_45.
- Wang ZY, Song J, Zhang DS. Nanosized As₂O₃/Fe₂O₃ complexes combined with magnetic fluid hyperthermia selectively target liver cancer cells. *World J Gastroenterol*. 2009;15(24):2995–3002. <https://doi.org/10.3748/wjg.15.2995>.
- Dokmanovic M, Marks PA. Prospects: histone deacetylase inhibitors. *J Cell Biochem*. 2005;96(2):293–304. <https://doi.org/10.1002/jcb.20532>.
- Daly K, Shirazi-Beechey SP. Microarray analysis of butyrate regulated genes in colonic epithelial cells. *DNA Cell Biol*. 2006;25(1):49–62. <https://doi.org/10.1089/dna.2006.25.49>.
- de Ruijter AJ, Meisma RJ, Bosma P, Kemp S, Caron HN, van Kuilenburg AB. Gene expression profiling in response to the histone deacetylase inhibitor BL1521 in neuroblastoma. *Exp Cell Res*. 2005;309(2):451–67. <https://doi.org/10.1016/j.yexcr.2005.06.024>.
- Peart MJ, Smyth GK, van Laar RK, Bowtell DD, Richon VM, Marks PA, et al. Identification and functional significance of genes regulated by structurally different histone deacetylase inhibitors. *Proc Natl Acad Sci U S A*. 2005;102(10):3697–702. <https://doi.org/10.1073/pnas.0500369102>.
- Rada-Iglesias A, Enroth S, Ameur A, Koch CM, Clelland GK, Respuela-Alonso P, et al. Butyrate mediates decrease of histone acetylation centered on transcription start sites and down-regulation of associated genes. *Genome Res*. 2007;17(6):708–19. <https://doi.org/10.1101/gr.5540007>.
- Goldstein JL, Brown MS. A century of cholesterol and coronaries: from plaques to genes to statins. *Cell*. 2015;161(1):161–72. <https://doi.org/10.1016/j.cell.2015.01.036>.
- Sun B, Jia YM, Hong J, Sun QW, Gao SX, Hu Y, et al. Sodium butyrate ameliorates high-fat-diet-induced non-alcoholic fatty liver disease through peroxisome proliferator-activated receptor α -mediated activation of β oxidation and suppression of inflammation. *J Agric Food Chem*. 2018;66(29):7633–42. <https://doi.org/10.1021/acs.jafc.8b01189>.
- Aguilar EC, da Silva JF, Navia-Pelaez JM, Leonel AJ, Lopes LG, Menezes-Garcia Z, et al. Sodium butyrate modulates adipocyte expansion, adipogenesis, and insulin receptor signaling by upregulation of PPAR- γ in obese Apo E knockout mice. *Nutrition*. 2018;47:75–82. <https://doi.org/10.1016/j.nut.2017.10.007>.

37. Wu SP, Zhang YG, Lu R, Xia YL, Zhou D, Petrof EO, et al. Intestinal epithelial vitamin D receptor deletion leads to defective autophagy in colitis. *Gut*. 2015;64(7):1082–94. <https://doi.org/10.1136/gutjnl-2014-307436>.
38. Sałkowska A, Karaś K, Walczak-Drzewiecka A, Dastych J, Ratajewski M. Differentiation stage-specific effect of histone deacetylase inhibitors on the expression of ROR γ T in human lymphocytes. *J Leukoc Biol*. 2017;102(6):1487–95. <https://doi.org/10.1189/jlb.6A0617-217R>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

